

Site-specific Glycosylation of Secretory Immunoglobulin A from Human Colostrum

Jincui Huang, Andrés Guerrero, Evan Parker, John S Strum, Jennifer T. Smilowitz, J. Bruce German, and Carlito B. Lebrilla

J. Proteome Res., **Just Accepted Manuscript** • DOI: 10.1021/pr500826q • Publication Date (Web): 28 Jan 2015

Downloaded from <http://pubs.acs.org> on February 6, 2015

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



Site-specific Glycosylation of Secretory Immunoglobulin A from Human Colostrum

Jincui Huang¹, Andres Guerrero¹, Evan Parker¹, John S. Strum¹, Jennifer T. Smilowitz^{2,3}, J. Bruce
German^{2,3} and Carlito B. Lebrilla^{1,4*}

1. Department of Chemistry, University of California, Davis, CA 95616, USA.
2. Foods for Health Institute, University of California, Davis, CA 95616, USA.
3. Department of Food Science and Technology, University of California, Davis, CA 95616, USA.
4. Department of Biochemistry and Molecular Medicine, University of California, Davis, CA 95616, USA.

Keywords:

Glycoproteomics, Glycan microheterogeneity, Mass spectrometry, sIgA, Site-specific glycosylation.

To whom correspondence should be addressed:

Carlito B. Lebrilla

E-mail: cblebrilla@ucdavis.edu.

Tel: 01-530-752-5054.

Fax: 01-530-752-8995.

Abbreviations

sIgA: secretory immunoglobulin A

pIgR: polymeric immunoglobulin receptor

HPLC: high-performance liquid chromatography

MFE: molecular feature extractor

Q-TOF: quadrupole time-of-flight

CID: collision-induced dissociation

SPE: solid phase extraction

Hex: hexose

HexNAc: N-acetylhexosamine

Neu5Ac: neuraminic acid (sialic acid)

Fuc: fucose

GCC: graphitized carbon cartridge

ACN: acetonitrile

HR: hinge region

ECC: Extracted compound chromatogram

ABSTRACT

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Secretory Immunoglobulin A (sIgA) is a major glycoprotein in milk and plays a key role in mediating immune protection of the gut mucosa. Although it is a highly glycosylated protein, its site-specific glycosylation and associated glycan micro-heterogeneity has still not been fully elucidated. In this study, the site-specific glycosylation of sIgA isolated from human colostrum (n = 3) was analyzed using a combination of LC/MS and LC/MS/MS and in-house software (Glycopeptide Finder). The majority of the glycans found are bi-antennary structures with one or more acidic Neu5Ac residue, however a large fraction belonged to truncated complex structures with terminal GlcNAc. Multiple glycosites were identified with nearly 30 glycan compositions located at seven sites on the secretory component, six compositions at a single site on the J-chain, and 16 compositions at five sites on the IgA heavy (H) chain. Site-specific heterogeneity and relative quantitation of each composition and the extent of occupation at each site was determined using non-specific proteases. Additionally, 54 O-linked glycan compositions located at the IgA1 hinge region (HR) were identified by comparison against a theoretical O-glycopeptide library. This represents the most comprehensive report to date detailing the complexity of glycan micro-heterogeneity with relative quantitation of glycoforms for each glycosylation site on milk sIgA. This strategy further provides a general method for determining site-specific glycosylation in large protein complexes.

INTRODUCTION

Secretory immunoglobulin A (sIgA) is the dominant immunoglobulin in the gut mucosal surface giving the mucosa specialized innate and adaptive defense against ingested pathogens and their toxins¹⁻³. sIgA is a protein complex consisting of two identical IgA monomers, joined together by a 16-kDa joint chain (Jchain) with one potential N-glycosylation site, and the secretory component (SC) with seven potential N-glycosylation sites⁴⁻⁶. The four polypeptide chains of sIgA are produced by two distinct cell types⁶. Plasma cells close to the epithelium produce dimeric IgA with an attached J chain, while epithelial cells express the polymeric immunoglobulin receptor (pIgR) that binds to dimeric IgA. Notably, the SC is derived from the pIgR after cleavage of the trans-membrane tail on the surface of epithelial cells⁷.

Glycosylation is one of the most common but also the most structurally complicated post-translational modification (PTM) of proteins. Approximately 70% of human proteins are predicted to be glycosylated^{8, 9}. sIgA is heavily glycosylated with both N- and/or O-linked oligosaccharides¹⁰, which vary according to the isotype and allotype of the immunoglobulin. Its oligosaccharides are attributed significant roles in immune protection, cell signaling, cell-cell recognition, microbial adhesion and invasion¹¹⁻¹³. sIgA is one of the major antibodies produced in human milk and is essential for the passive immunity of infants against infection¹⁴⁻¹⁶. Specific binding between sialylated glycans of sIgA and pathogens such as S-fimbriated *Escherichia coli* protects newborns from sepsis and meningitis due to infection¹⁷. The large diversity of N-glycan structures particularly on the SC provides many glycan epitopes that are potent decoys for lectins on bacterial surfaces thereby inhibiting infection by attachment with epithelial surfaces¹⁸⁻²⁰. sIgA also carries galactose-terminating glycans that protect against toxins including ricin, a particularly toxic galactose-specific lectin²¹.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Despite numerous studies establishing the effective protection provided by sIgA, the glycan heterogeneity and glycosite occupancy of this highly glycosylated protein remains incomplete. There have been studies showing the glycosylation of IgA in serum^{22, 23}, however milk is unique in that the protein also contains a highly glycosylated secretory component¹⁰. The first comprehensive attempt to characterize the glycosylation of sIgA from a commercial sample was reported by Royle *et al.*⁴, whereby polypeptide components were separated using gel-electrophoresis. Glycans were released for the respective gel spots corresponding to different polypeptides and characterized by mass spectrometry. The nature of the method yielded no site-specific information. Furthermore, the glycoforms spread the location of the glycoprotein on the gel, so releasing glycans from the largely-spread gel spot may not necessarily represent the majority of the glycoforms²⁴. Site-specific N-glycan analysis of sIgA was first performed by Deshpande *et al.* using in-gel trypsin digestion of a commercial sample²⁵. The results provided the characterization of several sites with the associated glycan heterogeneity. However, trypsin often produces large peptides, and in this case it could not resolve the glycosylation of ⁸³Asn and ⁹⁰Asn on the secretory component. Additionally, missed cleavages are common occurrences particularly if the tryptic sites are near sites of glycosylation²⁶⁻²⁸. For these reasons, employing only tryptic digestion will often yield incomplete glycan information.

A more complicated aspect is the O-glycosylation analysis of sIgA. The IgA1 hinge region contains up to nine potential O-glycosylation sites clustering in a 19 amino acid peptide sequence²⁹. A variety of MS analytical approaches have been used to characterize the glycan heterogeneity present in this portion of the immunoglobulin with only moderate success³⁰⁻³³. Additionally, IgA1 O-glycan site localization has been reported combining different protease digestions and

1
2
3 ECD/ETD^{29, 32, 33}. However in these studies sialic acid was removed enzymatically in order to
4
5
6 simplify the glycopeptide heterogeneity.
7

8
9 In this paper, we present an extensive and comprehensive glycan analysis of sIgA
10
11 from commercial pooled human colostrum and isolated from milk colostrum collected from three
12
13 healthy women. This laboratory has developed methods for site-specific analysis using
14
15 combinations of specific and non-specific proteolysis³⁴⁻⁴⁰. The method includes an automated
16
17 annotation software (Glycopeptide Finder) to analyze data from a nanoflow liquid
18
19 chromatography coupled with quadrupole time-of-flight analyzer (nanoLC-Q-TOF). This
20
21 analysis yielded both N- and O-glycosylation. Glycopeptide assignments were based on a
22
23 combination of accurate mass, retention time analysis and tandem MS yielding glycopeptides
24
25 with scalable lengths but including both non-specific and specific cleavages.
26
27
28
29
30
31
32

33 **Materials and Methods**

34 *Chemicals and Materials*

35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
SSL7/Agarose and phosphate-buffered saline (PBS), and TRIzol were purchased from
Invitrogen (San Diego, CA). Pronase E, cyanogen bromide (CNBr) activated sepharose 4B (S4B)
beads, and pooled human colostrum IgA were purchased from Sigma-Aldrich (St. Louis, MO).
PNGase F was purchased from New England Biolab (Ipswich, MA). Sequencing grade modified
trypsin and proteoMAX solution were purchased from Promega (Madison, WI). Dialysis tubing
(10MWCO) were purchased from Spectrum Lab (Rancho Dominguez, CA). Ammonium acetate
and acetic acid were of analytical grade from Merck (Darmstadt, Germany). Graphitized carbon
cartridges were purchased from Grace Davison Discovery Sciences (Deerfield, IL).

Human Colostrum Samples

Colostrum samples were collected from three healthy donors enrolled in the UC Davis Lactation Study who gave birth to term infants (> 38 weeks). Colostrum samples were manually collected on day 3-6 postpartum from one breast and transferred into polypropylene Falcon tubes, and frozen immediately in their kitchen freezers (-20 °C) until weekly sample pick up by the study staff. Samples were transported to the lab on dry ice and stored in -80 °C until processing.

The University of California Davis Institutional Review Board approved all aspects of the study and informed consent was obtained from all subjects. This trial was registered on clinicaltrials.gov (ID: NCT01817127).

Secretory Immunoglobulin A Extraction

sIgA from donor colostrum was extracted using Staphylococcal Superantigen-Like Protein 7 (SSL 7)/agarose⁴¹. Briefly, whole human colostrum samples (0.5 mL) were centrifuged at 4000 xg, for 30 minutes at 4°C. A CaCl₂ solution (pH 4.6) was added to the lower layer after the removal of top layer fat and lipids to a final concentration of 60 mM. The mixture was incubated 1 hour at room temperature (~25°C), and further centrifuged at 6750 xg for 30 minutes. The top layer was extracted as whey proteins. The empty column was packed with 1ml of SSL 7/agarose resin and equilibrated with 1X phosphate buffer saline (running buffer). The whey fractions were loaded onto the columns and the flow-through was collected and reloaded onto the column three times. Upon washing with 10 mL of the running buffer, sIgA bound to SSL 7 was eluted with 5 mL of 0.1 M glycine (pH 2-3) followed by the pH adjustment of the eluate to pH 7.5 by adding a neutralization buffer (1M Tris pH 8). Fractions were collected, dialyzed using a dialysis membrane with a molecular weight cut-off of 10,000 against nano-pure water, concentrated and stored at -20°C. Protein concentration was determined using the Bradford assay and 5 µl aliquots

1
2
3 were assayed by SDS-PAGE to confirm protein purity. A 5- μ L sample was mixed with 5 μ L of
4
5 laemli buffer, 1 μ L of 550 mM dithiothreitol (DTT) at 60 °C for 1 hour prior to incubating with 2
6
7 μ L of 450 mM iodoacetamide (IAA) for 30 min in the dark. The SDS-PAGE was performed at
8
9 140 V for 1 hr, followed by staining with coomassie blue for 1 hr. The gel was then unstained
10
11 with water overnight.
12
13

14 15 *Trypsin Digestion* 16 17

18
19 A 100 μ L aliquot of 50 mM NH_4HCO_3 buffer was added to 10 μ g isolated sIgA as described
20
21 above, followed by reducing with 2 μ L of 550 mM DTT for 50 min at 60 °C. A 4 μ L solution of
22
23 450 mM IAA was then added, and carboxymethylation was performed by incubation for 30 min
24
25 at room temperature in the dark. Reduced and carboxymethylated sIgA was digested using
26
27 trypsin 1 μ L (1 μ g/ μ L) in 100 μ L of 50 mM NH_4HCO_3 buffer, for 18 hr at 37 °C. The digests
28
29 were purified on a reverse-phase C18 pipette zip-tip (Agilent Technology, Wilmington, DE,
30
31 USA). The C18 zip-tip was preconditioned successively with acetonitrile (ACN) (150 μ L, 10
32
33 times) and water (150 μ L, 10 times). The tryptic digest was loaded on to the zip-tip by pipetting
34
35 20 times followed by similar 10 washings with nanopure water (150 μ L). Tryptic peptides from
36
37 sIgA were eluted with 0.05% formic acid (FA) in 80% ACN in water (v/v) (200 μ L, 20 times
38
39 extractions), dried down, and reconstituted in 20 μ L nanopure water. Standard proteomic analysis
40
41 of the extracted sIgA was performed to further verify the purity.
42
43
44
45
46

47 48 *Pronase E Digestion and Glycopeptide Cleanup* 49

50
51 Pronase E was covalently coupled to CNBr activated sepharose beads via coupling chemistry
52
53 as reported previously^{34, 36, 38}. A 100- μ g sample was dissolved in 100 μ L of 50 mM NH_4HCO_3
54
55 buffer followed by reduction with 2 μ L of 550 mM DTT for 50 min at 60 °C. A 4- μ L solution of
56
57 450 mM IAA was added, and carboxymethylation was performed for 50 min at room temperature
58
59
60

1
2
3 in the dark. Reduced and carboxymethylated samples were added to the pronase beads and
4
5 incubated at 37 °C for 18 h with gentle agitation. The glycopeptide digest was desalted and
6
7 enriched via solid phase extraction (SPE) using graphitized carbon cartridges. After
8
9 loading the glycopeptide on preconditioned 150-mg bed volume graphitized carbon cartridges
10
11 (Grace Davison Discovery Sciences, Deerfield, IL, USA), the cartridges were washed with 6-mL
12
13 nanopure water. A clean mixture of glycopeptides were eluted with 6 mL of 0.05%
14
15 Trifluoroacetic acid (TFA) in 40% ACN in water (v/v) and 6 mL of 0.05% TFA in 80% ACN in
16
17 water (v/v). Collected fractions were dried down and then reconstituted in 20 µL of nanopure
18
19 water. A 2-µL volume was used for MS analysis.
20
21
22
23
24

25 *Determination of O-glycosylation by PNGase F followed by Trypsin Digestion*

26
27

28 A 10-µg sample of sIgA dissolved in 50 µL NH₄HCO₃ buffer solution was reduced by DTT (1 µL,
29
30 550 mM) and carboxymethylated by adding IAA (2 µL, 450 mM). The solution was digested by
31
32 the addition of PNGase F (1 µL) for 16 hours at 37 °C to remove the N-glycans. Trypsin
33
34 digestion was added to the solution, which was incubated at 37 °C for 18 hours for digestion.
35
36 Glycopeptides in the digest were purified via hydrophilic affinity separation according to a
37
38 method described previously⁴². Briefly, O-glycopeptides were enriched with a 15 µL packed
39
40 volume of sepharose CL4B in 1 mL of an organic solvent of 1-butanol/ethanol/H₂O (5:1:1, v/v).
41
42 Followed by shaking for 45 min, the gel was washed twice with the same organic solvent prior to
43
44 incubating with an aqueous solvent of ethanol/H₂O (1:1, v/v) for 30 min. The solution phase was
45
46 collected after centrifugation and dried prior to reconstitution with 20 µL nanopure water.
47
48
49
50

51 *LC/MS analysis of glycopeptides*

52
53

54 A nano-HPLC-Chip Q-TOF instrument using the Agilent 1200 series microwell-plate
55
56 autosampler (maintained at 6 °C by the thermostat), capillary pump, nano pump, HPLC-Chip
57
58
59
60

1
2
3 interface, and the Agilent 6520 Q-TOF MS (Agilent Technologies, Inc., Santa Clara, CA) was
4 used in this study. The chip is equipped with a 40 nL enrichment column and a 43 x 0.075 mm
5 ID analytical column. Agraphitized carbon stationary phase was used for pronase produced
6 glycopeptides, while C18 stationary phase was used for the tryptic peptides. Both mobile phases
7 consists of 0.1% formic acid in 3% ACN in water (v/v) and 0.1% formic acid in 90% ACN in
8 water(v/v) as solvent A and B, respectively. The gradient was performed on the analytical
9 column to separate the pronase glycopeptides with a flow rate at 0.4 $\mu\text{L}/\text{min}$. The samples were
10 eluted in 45 min with the following gradient: 0% B (0.00-2.50 min); 0 to 16% B (2.50-20.00
11 min); 16 to 44% B (20.00-30.00 min); 44 to 100% B (30.00-35.00 min) and 100% B (35.00 -
12 45.00 min). The tryptic glycopeptides/peptides were eluted in 87min with the following gradient:
13 1% to 8% B (0.00-5.00 min); 8% to 26.5% B (5.00 min-48.00 min); 26.5% to 73% B (48.00
14 min-75 min); 73% to 99% (75.00 min-77.00 min); 99%B (77.00 min-87.00 min).

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The Agilent 6520 Q-TOF MS was operated in the positive ion mode for MS and MS/MS analysis. The recorded mass ranges were m/z 500-3,000 for MS-only and m/z 50-3,000 for MS/MS. Acquisition rates were 0.63 spectrum/s for both MS and MS/MS. All mass spectra were internally calibrated using the G1969-85000 ESI tuning mix (Agilent Technologies, Inc., Santa Clara, CA), with reference masses at m/z 922.010, and 1,521.971 in the positive ion mode. The parameters used were as follows: drying gas (N_2) temperature 325°C, drying gas flow rate 3 L/min, capillary voltage 1800 V, fragmentor voltage 150 V, skimmer voltage 65 V, and capillary current 0.08 μA , respectively. For MS/MS mode, the collision energies for each compound were calculated as follows:

$$V_{\text{collision}} = 1.8V \left(\frac{m/z}{100\text{Da}} \right) - 2.4V$$

Data Processing

1
2
3 The MS/MS data of the tryptic peptides were analyzed using X! Tandem (www.thegpm.org). X!
4 Tandem was set up to search the Swissprot human complete proteome database. X! Tandem was
5 searched with a fragment ion mass tolerance of 80 ppm and a parent ion tolerance of 100 ppm.
6 Iodoacetamide derivative of cysteine was specified in X! Tandem as a fixed modification.
7 Deamidation of asparagine and glutamine, oxidation of methionine and tryptophan were
8 specified in X! Tandem as variable modifications.
9

10 Data analyses for N-glycopeptides were performed with the MassHunter Qualitative Analysis
11 software ver. B.03.01 (Agilent Technologies, Inc., Santa Clara, CA). Deconvolution was
12 performed in the software with the grouping peak spacing tolerant of 0.0025 m/z plus 7 ppm and
13 isotope mode was set for peptides. The mass list of the glycopeptide precursor ions and the
14 associated tandem spectra were analyzed via an in-house software (Glycopeptide Finder)³⁹ set to
15 a 5% false-discovery rate. All N-glycopeptide assignments were made based on the mass of each
16 potential glycopeptide, the amino acid sequence(s) of the protein(s), and a specific tolerance
17 level (< 20 ppm). The presence of carbohydrate-specific oxonium fragment ions was used as
18 diagnostic ions to sort the product tandem spectra.
19

20 For the O-glycopeptide identification, the tryptic digest after PNGase F treatment was
21 analyzed in both MS and MS/MS mode. A theoretical O-glycopeptide library was created using
22 the tryptic peptide from the HR (HYTNPSQDVTVPCVPSTPPTPSPSTPPTPSPSCCHPR) with
23 glycan compositions within a specified range (HexNAc 0-6, Hex 0-6, Sialic Acid 0-12) based on
24 previous results^{25, 30, 33} and 0 to 3 carbamidomethyl groups. The resulting library is composed by
25 780 theoretical HR O-glycopeptides ranging in mass between 3964.818 Da and 9819.821
26 Da being the smaller difference between compounds larger than 1 Da. The library included
27 empirical formulae in order to compare isotopic patterns. MS data was analyzed with
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 MassHunter Qualitative Analysis software ver. B.03.01 (Agilent Technologies, Inc., Santa Clara,
4 CA). Compound ion signals with different charge states were grouped using the Molecular
5 Feature algorithm(MFE) and compared against the O-glycopeptide library with an error
6 tolerance of 30 ppm. Only ions with a signal-to-noise ratio above 3 were considered. Isotopic
7 distributions were manually checked. The extracted compound chromatograms (ECCs) for each
8 O-glycopeptide were obtained and their corresponding retention times compared.
9
10
11
12
13
14
15
16
17
18
19

20 21 **Results and Discussion**

22 *Glycan-heterogeneity of sIgA from Human Colostrum*

23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Colostrum sIgA was extracted from three individual donors. Isolation of sIgA was performed
as outlined in the methods section. The purity of sIgA was assessed by SDS-PAGE with four
main bands corresponding to the SC, H chain and light chain of IgA, and J chain as shown in
Figure S1. Standard proteomic analysis was performed on the isolated sIgA to further confirm
the purity of the protein. The results are listed in **Table S1** and indicate that the isolated sIgA is
relatively pure with only very minor protein contaminants.

The extracted compound chromatograms (ECCs) produced by the molecular feature extractor
(MFE) demonstrate similar glycopeptide profiles as well as abundances for the three colostrum
sIgA samples (**Figure S2**). A list of glycopeptides from the pronase-digested sIgA isolated from
one representative colostrum sample is shown in **Table 1**. The ECCs produced by the MFE
yielded intensities that were used for obtaining relative abundances. Abundances for all peptides
and glycoforms representing a specific site were added³⁷.

The elution order of glycopeptides on PGC follows loosely a set of empirical rules^{43, 44}.
Smaller peptides elute before larger ones with the same glycoforms. For example, the most

1
2
3 abundant complex-type glycan, Hex₅HexNAc₄Fuc₁NeuAc₂ attached to the peptide
4
5 HVKHYT⁹²NPSQ from IgA2, eluted late at 33.9 min, while the same glycan composition on the
6
7 peptide ¹³⁵NDT from the SC region eluted much earlier at 17.8 min. Meanwhile, increasing the
8
9 number of sialic acid residues increases the elution time. For example, the glycopeptide
10
11 ⁷¹NISDPTSPL+Hex₅HexNAc₄NeuAc₁ (21.7 min) with one sialic acid eluted earlier than the
12
13 glycopeptide ⁷¹NISDPTSPL+Hex₅HexNAc₄NeuAc₂ (25.5 min) with two sialic acids. In general,
14
15 sialylated species elute later than neutral species^{43, 44}.
16
17
18
19

20 **Figure 1A** shows the overlaid ECCs of glycopeptides from isolated sIgA with the major peaks
21
22 annotated. The most abundant peak was determined to
23
24 beHVKHYT⁹²NPSQ+Hex₅HexNAc₄Fuc₁NeuAc₂ from the conserved site (⁹²Asn) of IgA2 (34
25
26 min). Glycosylation on this site is unusual for the reasons discussed below. More IgA2
27
28 glycopeptides were observed associated with complex sialylated or high mannose type glycans
29
30 (peaks in yellow). Glycopeptides corresponding to SC (peaks in green) were observed mostly
31
32 with complex type glycans varying from biantennary to tetraantennary with a varying number of
33
34 Neu5Ac residues eluting mostly from 16–21 minutes, and 23–29 min. The J chain glycopeptides
35
36 (peaks in pink) were among the less intense peaks and originated from the single site on the J
37
38 chain. All the glycopeptides from the J chain were observed to contain at least one Neu5Ac. The
39
40 glycopeptides eluting between 7 min to 29 min illustrate the large complexity and the dynamic
41
42 range of the glycosylation (**Figures 1B** and **1C**).
43
44
45
46
47

48 To obtain structural information from the peaks in **Figure 1**, tandem MS spectra were
49
50 performed (shown in **Figure S3**). We have previously reported characteristic fragmentation of
51
52 glycopeptides. Glycan fragmentation yielding B-type ions is commonly observed in the collision
53
54 induced dissociation (CID) spectra of glycopeptides⁴⁵. A typical tandem MS spectrum is shown
55
56
57
58
59
60

1
2
3 for the glycopeptide E⁷¹NISDPTSPL+Hex₅HexNAc₃Fuc₁NeuAc₁ (**Figure 2A**). The tandem mass
4
5 spectrum shows fragment products consistent with hybrid-type glycan structures. The fragments
6
7 include a series of Hex losses (162 Da) prior to the loss of core HexNAc (203 Da). Similarly, Y-
8
9 type ions corresponding to the initial neutral loss of (Hex+Fucose) are also associated with hybrid
10
11 structures. Core fucosylation was confirmed via the loss of fucose (146 Da) to yield the Y₁ ion of
12
13 (E⁷¹NISDPTSPL+HexNAc₁). **Figure 2B** is representative of deconvoluted MS/MS spectrum for
14
15 a SC glycopeptide containing a neutral complex type tetraantennary N-glycan
16
17 (VSLEVSQGPGLL¹³⁵NDTK+Hex₇HexNAc₆). In this study, higher abundances of
18
19 tetraantennary structures were detected from sIgA compared with earlier work ⁴. **Figure 2C**
20
21 depicts the CID data with the loss of the two Neu5Ac (291 Da) residues to yield the extent of
22
23 sialylation. With sialylation, B-type ions with neutral masses of 273 Da, 291 Da, 656 Da, and
24
25 818 Da were observed corresponding to Neu5Ac–H₂O, Neu5Ac, Neu5Ac+Hex+HexNAc, and
26
27 Neu5Ac+Hex₂+HexNAc. Due to the labile nature of sialic acids and the fact that they generally
28
29 reside in the terminal positions, losses of Neu5Ac and Neu5Ac+Hex+HexNAc from the
30
31 precursor peaks are common.
32
33
34
35
36
37

38
39 In several of the large glycopeptides, the loss of H₂O, specifically from Neu5Ac, was observed.
40
41 For example, after deconvolution, glycopeptides with neutral masses 4554.9 Da and 4572.9 Da,
42
43 differing by 18 Da, were assigned as LTNFPE⁹⁰NGTFVVNIAQLSQ+Hex₆HexNAc₆NeuAc₂–
44
45 H₂O and LTNFPE⁹⁰NGTFVVNIAQLSQ+Hex₆HexNAc₆NeuAc₂. **Figure S4** represents
46
47 deconvoluted MS/MS spectra for these two glycopeptides. Neutral loss of Neu5Ac (291 Da) was
48
49 observed followed by the initial loss of one Hex (**Figure S4A**). However, **Figure S4B** reveals a
50
51 loss of 273 Da corresponding to (Neu5Ac – H₂O) as the first loss. Y-type ions with the intact
52
53
54
55
56
57
58
59
60

1
2
3 peptide, LTNFPE⁹⁰NGTFVVNIAQLSQ associated with the trimannosyl core are present in both
4
5 spectra indicating the same peptide were present in the two glycopeptides.
6
7
8
9

10 *Glycan-heterogeneity of pooled human colostrums IgA*

11
12 The same experiments were performed on pooled human colostrum sIgA obtained
13 commercially. A list of glycopeptides from the pronase-digested sIgA is shown in **Table S2**.
14
15 A much lower diversity of peptides and glycan structures were observed with the commercial
16
17 pooled sIgA as compared to the isolated sIgA from individual donors. The glycopeptide
18
19 HVKHYT⁹²NPSQ+Hex₅HexNAc₄Fuc₁NeuAc₂ from sIgA H chain remained the most abundant
20
21 species. However, ¹⁸⁶Asn was not found with glycosylation in the pooled sIgA. In general, sIgA
22
23 extracted from the colostrum from individuals contained greater glycan diversity than those
24
25 obtained from the commercial sIgA, including more triantennary and tetraantennary glycan
26
27 structures at multiple sites particularly in the SC.
28
29
30
31
32
33

34 **Figure 3** shows the overlaid ECCs of the glycopeptides from the commercial sIgA sample
35
36 with the major peaks annotated. The majority of the observed glycopeptides contained Neu5Ac
37
38 residue(s) with complex-type bi-antennary N-glycans. Included in this group are several truncated
39
40 structures with terminal GlcNAc. Although less abundant, other glycan types were observed,
41
42 including high mannose and hybrid. Over 80% of the glycopeptide abundances belonged to
43
44 structures with core fucosylation, which were verified by tandem MS (see below). In general,
45
46 essentially all singly fucosylated glycans were core fucosylated. Only one glycan was observed
47
48 with two fucose residues.
49
50
51
52

53 **Figure S5 and S6** show the deconvoluted MS/MS spectra of several N-glycopeptides
54
55 representing different sites on the pooled sIgA. **Figure S5A and S5B** are representative of two
56
57
58
59
60

1
2
3 deconvoluted MS/MS spectra of the glycopeptides from J chain and SC, respectively. The
4
5 fragment denoted as peptide+HexNAc was observed and highly informative for identifying the
6
7 two glycopeptides. **Figure S6A** shows the fragmentation of one of the most abundant
8
9 glycopeptide(HVKHYT⁹²NPSQ+Hex₅HexNAc₄Fuc₁NeuAc₂). This peptide is highly unusual
10
11 because it does not conform to the N-X-S sequon²⁵, where X cannot be proline. However, we
12
13 have a number of confirming evidences for the glycosylation of this site. The tandem MS shows
14
15 the progression of saccharide loss to the peptide+GlcNAc core (1412.65 Da), which gives
16
17 indication as to the mass of the peptide. The GP Finder software does not provide other
18
19 structures corresponding to this mass that were consistent with the tandem MS. Additionally,
20
21 tandem MS of other glycopeptides representing the same site was also obtained. For example,
22
23 tandem MS spectra representative for glycopeptide YT⁹²NPSQDV+Hex₅HexNAc₄NeuAc₁ and
24
25 glycopeptide YT⁹²NPSQDV+ Hex₅HexNAc₄NeuAc₂are shown in **Figure S6B** and **S6C**, which
26
27 provides the glycan heterogeneity of this site. The fact that glycan heterogeneity was observed at
28
29 this site with various peptide backbone as shown in **Table 1** is a confirmation that this site is
30
31 glycosylated. There are also reports in literature that support glycosylation on this site. In a
32
33 report by Picariello *et al.*, they found tryptic peptides from this site is potentially modified and
34
35 glycosylated⁴⁶.A 1Da mass shift from the peptide after PNGase F release is indicative of a
36
37 putative N-glycan site.

38
39
40
41
42
43
44
45
46 Assignment of the glycopeptide structures using the software GP Finder was performed with
47
48 95% confidence based on decoy data. Some assignments were complicated by the large number
49
50 of potential sites for sIgA. One complication is when the peptide mass was indistinguishable
51
52 between different glycopeptides from different sites. For example, the glycopeptide with the
53
54 peptide sequence E⁹⁰NGT from the SC and SEA^{131/144}N from IgA1/IgA2 have nearly identical
55
56
57
58
59
60

1
2
3 masses corresponding to 419.165 Da. Tandem MS could not resolve the peptides because the
4 peptides were short and did not yield suitable sequence data. Other problems arose when the
5 peptides were even shorter. For example peptides such as NV and AN were observed as
6 products and corresponded to four different glycosylation sites. It is noted that these
7 glycopeptide abundances would affect the determination of site occupancy. However, these short
8 peptides were minor products with very low abundances (<5%).

9
10 Compared with previous glycan analyses of sIgA^{4, 23, 25, 46}, our method revealed a more
11 comprehensive glycan micro-heterogeneity from each site. We find glycans with heterogeneity
12 for all potential N-sites including seven for the SC, two for IgA1, five for IgA2, and one for the J
13 chain. The most comprehensive analysis before this report obtained five sites from the SC, two
14 sites for IgA1, two IgA2 and one for the J chain²⁵. A potentially controversial assignment is that
15 for ⁹²Asn on IgA2, because it goes against the putative sequon. We find this site to be
16 glycosylated, consistent with an earlier proposal but not found in a subsequent analysis²⁵.

36 *Semi-quantitative Site-specific Glycosylation*

37
38 A summary of detailed and comprehensive glycan site heterogeneity of the sIgA isolated from
39 a single human colostrum sample is depicted in **Figure 4A-C**. The result of the pooled
40 commercial sample is shown in **Figure 4D-F**. The sIgA glycopeptide profiles were found to be
41 similar between the fresh human colostrum samples from three donors. However, these
42 samples were more highly glycosylated than the pooled commercial sample. For example,
43 commercial sIgA was not found glycosylated in ¹⁸⁶N of the SC while fresh colostrum was
44 glycosylated. A greater site-heterogeneity was observed with most of the sites in the fresh
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 colostrum sIgA. The result is not unexpected as commercial enrichment, chemical processing,
4 and the storage may degrade glycans.
5
6

7
8 Quantitative information was obtained based on the abundances of the glycopeptides. Two
9 types of quantitative information are provided in the figures. The small circular symbol under the
10 glycan structure designates the abundance of that glycan relative to all the glycans at that
11 glycosite. The circular symbol under the amino acid represents the occupancy of the site. To
12 obtain these values, the MS abundances for all glycoforms associated with a site were added.
13 The abundances were then combined for each glycoform and normalized to the total site
14 abundance for the relative glycan abundance at each site. The sum for each site represents the
15 occupancy while the sum for each glycan represents the relative abundance of the glycoform.
16 The occupancy is further normalized to the most occupied site (100%). Quantitation of glycan
17 site occupancy remains a difficult task. We propose a semi-quantitative method using ion
18 abundances but, for the moment, neglecting issues such as differences in ionization efficiency of
19 different peptides and different glycoforms as well as the dispersion of ion abundances across
20 different peptides due to the nature of the nonspecific proteases. Nonetheless, a previous study
21 by Hua *et al.* reported the quantitative accuracy and reproducibility of non-specific glycosylation
22 analysis and revealed the efficiency of this method⁴⁷. In this way, we find that the most
23 abundant site is ⁹²Asn of the IgA2 in the protein complex.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45
46 **Figure 4A** illustrates the site heterogeneity of the SC. The most abundant glycans at ¹³⁵Asn
47 correspond to the tetra-antennary compositions. The glycans on the SC illustrate general
48 diversity of the structures. The majority of structures were tri- and tetraantennary complex-type
49 glycans further decorated with Neu5Ac and core fucosylation, in accordance with previous
50 reports^{4, 25}. A small amount of high mannose (Man5 and Man6) and neutral complex glycan
51
52
53
54
55
56
57
58
59
60

1
2
3 structures were observed. Glycosylation was found to be most abundant at ⁹⁰Asn among the
4
5 seven sites on SC, followed by ⁸³Asn and ¹³⁵Asn.
6
7

8 The characterization of the site heterogeneity on the J chain is summarized in **Figure 4B**.
9
10 Krugman *et al.*⁴⁸ previously reported that the substitution of the Asn with Ala in J chain prevents
11
12 IgA dimer formation, indicating the significance of glycosylation to the tertiary structure. Also,
13
14 the glycans associated with J chain are essential for binding and trans-locating
15
16 pIgR across epithelial cells⁴⁹. However, the results obtained here show a single site (⁷¹Asn) with
17
18 less than 20% occupancy corresponding to mainly sialylated complex type glycans. The results
19
20 are consistent with previous work but it calls to question contribution of this glycosite given the
21
22 relatively low occupancy.
23
24
25

26
27 The H chain of IgA is significantly more glycosylated than the J chain and the SC. **Figure 4C**
28
29 details the abundances of site-specific IgA1 H chain and IgA2 H chain glycans. Because the
30
31 amino acid sequence near sites ¹⁴⁴Asn for IgA1 are identical to those of ¹³¹Asn for IgA2, and
32
33 similarly for ³⁴⁰Asn of IgA1 and ³²⁷Asn of IgA2, the glycopeptide assignments could not be more
34
35 precise and were therefore combined for both IgA1 and IgA2. The analysis shows that the
36
37 majority of N-glycans on the H chains are sialylated and/or terminated by galactose (Gal)
38
39 residues, in agreement with previous studies of serum IgA^{22, 50}. The results are therefore consistent
40
41 with the formation of milk sIgA, which originates from B cells. While ^{131/144}Asn and ^{327/340}Asn
42
43 sites were observed as being slightly glycosylated, the ⁹²Asn from IgA2 was characterized as
44
45 being mostly glycosylated mainly due to the most abundant glycopeptide
46
47 (HVKHYT⁹²NPSQ+Hex₅HexNAc₄Fuc₁NeuAc₂). Moreover, bi-antennary complex type glycans
48
49 were observed at ⁹²Asn associated with one or two sialic acids. Core fucosylation was detected,
50
51
52
53
54
55
56
57
58
59
60

1
2
3 as discussed previously from the MS/MS data, mainly due to the observation of the loss of
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

as discussed previously from the MS/MS data, mainly due to the observation of the loss of
fucose to yield an informative Y_1 ion.

Overall, from the colostrum of individual donors, sialylated glycans were found to be the most
abundant (82%), from which mono-sialylation (35%) and bi-sialylation (47%) were both present.
Fucosylation (36%) were observed mostly as core fucosylation.

In the commercial sIgA, the most abundant glycopeptide was
HVKHYT⁹²NPSQ+Hex₅HexNAc₄Fuc₁NeuAc₂ with ⁹²Asn being the most occupied site. The
absence of branched complex type N-glycans, in particular from the SC, was notable possibly
due to the processing methods of the commercial sample (**Figure 4D-4F**).

O-glycosylation Analysis

We have previously determined N- and O-glycosylation simultaneously for other
glycoproteins,^{24, 36, 37, 47} however the same comprehensive analysis described above failed to
detect O-glycopeptides in the hinge region of IgA. The presence of peptides and N-glycopeptides
in the IgA hydrolysate complicate the detection of the O-glycopeptides. In order to observe the
O-glycopeptides, we removed the N-glycans from IgA, digest the protein with trypsin, and
enriched the O-glycopeptides using hydrophilic affinity methods.

The enriched O-glycopeptide fraction was analyzed in both MS and MS/MS mode.
A glycopeptide library was employed to identify the glycopeptides using accurate mass and the
theoretical isotopic distribution. This approach revealed 54 compounds compiled in **Table 2**.
Complete cysteine alkylation was observed for all hinge region O-glycopeptides. The identified
compounds eluted in a narrow interval (between minute 22 and 28) in agreement with the elution
behavior in C18 of a family of glycopeptides sharing the same peptide moiety and multiple

1
2
3 glycan compositions. Tandem-MS of the O-glycopeptides (**Figure 5A-5C**) revealed
4 characteristic oxonium ions and the presence of peptide backbone fragment ion b11 (621.79 m/z
5 and 612.79 m/z) corresponding to the sequence HYTNPSQDVTV, which unambiguously
6 identifies the compounds as HR glycopeptides. Due to the overlap of the glycopeptide peaks and
7 their low abundances, tandem-MS spectra were obtained only for the more abundant species.
8 However, O-glycopeptide assignments were confirmed by comparing the elution order. The elution
9 order follows a certain pattern as illustrated in **Figure 6** for the O-glycopeptide series containing
10 five N-acetylgalactosamine, three galactose and different number of sialic acid residues.
11
12
13
14
15
16
17
18
19
20
21
22

23 In every case, the addition of sialic acid residues increases the retention time of the
24 corresponding O-glycopeptides (**Figure S7**). Similarly, the elution time order can be
25 compared with the number of hexoses or hexosamines residues. The addition of these
26 monosaccharides decreased gradually the retention time (**Figure S8**). The combined comparison
27 of the elution orders confirmed the compound assignments. Isomeric o-glycopeptides were found
28 for 5 different compositions when examining the extracted compounds chromatograms (**Figure**
29 **S7**). The results are in agreement with those published previously²⁵.
30
31
32
33
34
35
36
37
38
39

40 CONCLUSION

41
42
43 The results demonstrate the detection and characterization of a complicated large glycoprotein
44 complex IgA containing multiple polypeptides and numerous sites of glycosylation. N-
45 glycosylation analysis of sIgA from human colostrum is performed at the glycopeptide level
46 using pronase digestion and subsequent LC-MS and LC-MS/MS analysis. Extensive and
47 comprehensive site-specific glycosylation analysis was obtained, which will aid our understanding
48 of the complex biological roles of sIgA. O-glycosylation was identified via trypsin digestion with
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 54 most likely glycan compositions present on this multiply O-glycosylated peptide. This study
4
5 demonstrates the wealth of quantitative site-occupancy information on protein complexes that
6
7 include with multiple glycosites.
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Supporting Information

Table S1. Associated proteins identified in extracted colostrum sIgA

Table S2. Glycopeptides identified from pooled commercial sIgA

Figure S1. sIgA (5 $\mu\text{g}/\text{lane}$) and commercial available sIgA (20 $\mu\text{g}/\text{lane}$) was reduced, alkylated, and then run on a 10% Tris-HCl gel (from left to right are colostrum, transitional, mature and commercial sIgA)

Figure S2. Overlaid Extracted Compound chromatogram (ECC) in different colors from pronase digested sIgA isolated from three individuals.

Figure S3. MS/MS spectra used for assignment of glycopeptides. Notably, diagnostic ions were found in each spectrum corresponding to deconvoluted neutral mass of HexNAc (203.08 Da), Hex+HexNAc (365.13 Da), and Neu5Ac (291.09) if sialylated.

Figure S4. (A and B) Deconvoluted MS/MS spectra of two glycopeptides from SC with the same assignments. Green Circles, yellow circles, blue squares, red triangles, and purple diamonds represent mannose, galactose, GlcNAc, fucose and Neu5Ac, respectively.

Figure S5. Deconvoluted MS/MS spectra of two N-linked glycopeptides from sIgA. (A) MS/MS data for a N-linked peptide from J chain; (B) MS/MS data for a N-linked peptide from SC. Green Circles, yellow circles, blue squares, red triangles, and purple diamonds represent mannose, galactose, GlcNAc, fucose and Neu5Ac, respectively.

Figure S6. Deconvoluted MS/MS spectra of three N-linked glycopeptides for ^{92}ASN from IgA2. (A) MS/MS data for $\text{HVKHYT}^{92}\text{NPSQ}+\text{Hex}_5\text{HexNAc}_4\text{Fuc}_1\text{NeuAc}_2$; (B) MS/MS data for

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

YT⁹²NPSQDV+Hex₅HexNAc₄NeuAc₁; (C) MS/MS data for YT⁹²NPSQDV+Hex₅HexNAc₄NeuAc₂. Green Circles, yellow circles, blue squares, red triangles, and purple diamonds represent mannose, galactose, GlcNAc, fucose and Neu5Ac, respectively.

Figure S7. Extracted compound chromatograms (ECCs) of HR O-glycopeptides containing same number of N-acetylgalactosamine and galactose residues and a variable number of sialic acid monosaccharides. Isomeric compounds (*).

Figure S8: Retention time against number of monosaccharide residues for the different HR O-glycopeptide series.

Supporting Information, this material is available free of charge via <http://pubs.acs.org>.

Acknowledgement

The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Research reported in this publication was supported by the National Institute of Child Health and Human Development, National Institute of General Medicine, National Center of Complimentary and Alternative Medicine of the National Institutes of Health under award number R01HD061923, R01GM049077, and 1U24DK097154.

References

1. Renegar, K.; Small Jr, P., Passive immunization: systemic and mucosal. *Mucosal Immunology* **1994**, 3, 841-851.
2. Russell, M.; Kilian, M.; Lamm, M., Biological activities of IgA. *Mucosal immunology* **1999**, 225.
3. Phalipon, A.; Cardona, A.; Kraehenbuhl, J.-P.; Edelman, L.; Sansonetti, P. J.; Corthésy, B., Secretory component: a new role in secretory IgA-mediated immune exclusion in vivo. *Immunity* **2002**, 17, (1), 107-115.
4. Royle, L.; Roos, A.; Harvey, D. J.; Wormald, M. R.; Van Gijlswijk-Janssen, D.; Redwan, E.-R. M.; Wilson, I. A.; Daha, M. R.; Dwek, R. A.; Rudd, P. M., Secretory IgA N-and O-glycans provide a link between the innate and adaptive immune systems. *Journal of Biological Chemistry* **2003**, 278, (22), 20140-20153.
5. Johansen, F.-E.; Braathen, R.; Brandtzaeg, P., The J chain is essential for polymeric Ig receptor-mediated epithelial transport of IgA. *The Journal of Immunology* **2001**, 167, (9), 5185-5192.
6. Mostov, K. E., Transepithelial transport of immunoglobulins. *Annual review of immunology* **1994**, 12, (1), 63-84.
7. Underdown, B. J.; Mestecky, J., Mucosal immunoglobulins. *Handbook of mucosal immunology* **1994**, 79.
8. An, H. J.; Peavy, T. R.; Hedrick, J. L.; Lebrilla, C. B., Determination of N-glycosylation sites and site heterogeneity in glycoproteins. *Analytical chemistry* **2003**, 75, (20), 5628-5637.
9. Apweiler, R.; Hermjakob, H.; Sharon, N., On the frequency of protein glycosylation, as deduced from analysis of the SWISS-PROT database. *Biochimica et Biophysica Acta (BBA)-General Subjects* **1999**, 1473, (1), 4-8.
10. Zauner, G.; Selman, M. H.; Bondt, A.; Rombouts, Y.; Blank, D.; Deelder, A. M.; Wuhrer, M., Glycoproteomic analysis of antibodies. *Molecular & Cellular Proteomics* **2013**, 12, (4), 856-865.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
11. Takahashi, M.; Kuroki, Y.; Ohtsubo, K.; Taniguchi, N., Core fucose and bisecting GlcNAc, the direct modifiers of the N-glycan core: their functions and target proteins. *Carbohydrate research* **2009**, 344, (12), 1387-1390.
 12. Marth, J. D.; Grewal, P. K., Mammalian glycosylation in immunity. *Nature Reviews Immunology* **2008**, 8, (11), 874-887.
 13. Gu, J.; Sato, Y.; Kariya, Y.; Isaji, T.; Taniguchi, N.; Fukuda, T., A Mutual Regulation between Cell-Cell Adhesion and N-Glycosylation: Implication of the Bisecting GlcNAc for Biological Functions. *Journal of proteome research* **2008**, 8, (2), 431-435.
 14. Newburg, D. S.; Ruiz-Palacios, G. M.; Morrow, A. L., Human milk glycans protect infants against enteric pathogens. *Annu. Rev. Nutr.* **2005**, 25, 37-58.
 15. Hanson, L. A., Breastfeeding provides passive and likely long-lasting active immunity. *Annals of Allergy, Asthma & Immunology* **1998**, 81, (6), 523-537.
 16. Peterson, R.; Cheah, W. Y.; Grinyer, J.; Packer, N., Glycoconjugates in human milk: Protecting infants from disease. *Glycobiology* **2013**, 23, (12), 1425-1438.
 17. Schroten, H.; Stapper, C.; Plogmann, R.; Köhler, H.; Hacker, J.; Hanisch, F.-G., Fab-independent antiadhesion effects of secretory immunoglobulin A on S-fimbriated Escherichia coli are mediated by sialyloligosaccharides. *Infection and immunity* **1998**, 66, (8), 3971-3973.
 18. Wold, A. E.; Mestecky, J.; Tomana, M.; Kobata, A.; Ohbayashi, H.; Endo, T.; Edén, C. S., Secretory immunoglobulin A carries oligosaccharide receptors for Escherichia coli type 1 fimbrial lectin. *Infection and immunity* **1990**, 58, (9), 3073-3077.
 19. Falk, P.; Roth, K. A.; Boren, T.; Westblom, T. U.; Gordon, J. I.; Normark, S., An in vitro adherence assay reveals that Helicobacter pylori exhibits cell lineage-specific tropism in the human gastric epithelium. *Proceedings of the National Academy of Sciences* **1993**, 90, (5), 2035-2039.

- 1
2
3 20. Boren, T.; Falk, P.; Roth, K. A.; Larson, G.; Normark, S., Attachment of *Helicobacter pylori* to
4 human gastric epithelium mediated by blood group antigens. *Science* **1993**, 262, (5141), 1892-1895.
5
6
7
8 21. Mantis, N. J.; Farrant, S. A.; Mehta, S., Oligosaccharide side chains on human secretory IgA serve
9 as receptors for ricin. *The Journal of Immunology* **2004**, 172, (11), 6838-6845.
10
11
12 22. Tanaka, A.; Iwase, H.; Hiki, Y.; Kokubo, T.; Ishii-Karakasa, I.; Toma, K.; Kobayashi, Y.; Hotta, K.,
13 Evidence for a site-specific fucosylation of N-linked oligosaccharide of immunoglobulin A1 from normal
14 human serum. *Glycoconjugate journal* **1998**, 15, (10), 995-1000.
15
16
17 23. Gomes, M. M.; Wall, S. B.; Takahashi, K.; Novak, J.; Renfrow, M. B.; Herr, A. B., Analysis of IgA1
18 N-Glycosylation and Its Contribution to Fc α RI Binding[†]. *Biochemistry* **2008**, 47, (43), 11285-11299.
19
20
21 24. Nwosu, C. C.; Huang, J.; Aldredge, D. L.; Strum, J. S.; Hua, S.; Seipert, R. R.; Lebrilla, C. B., In-gel
22 nonspecific proteolysis for elucidating glycoproteins: a method for targeted protein-specific
23 glycosylation analysis in complex protein mixtures. *Analytical chemistry* **2012**, 85, (2), 956-963.
24
25
26 25. Deshpande, N.; Jensen, P. H.; Packer, N. H.; Kolarich, D., GlycoSpectrumScan: fishing
27 glycopeptides from MS spectra of protease digests of human colostrum sIgA. *Journal of proteome*
28 *research* **2010**, 9, (2), 1063-1075.
29
30
31 26. LOO, T. W.; CLARKE, D. M., The human multidrug resistance P-glycoprotein is inactive when its
32 maturation is inhibited: potential for a role in cancer chemotherapy. *The FASEB journal* **1999**, 13, (13),
33 1724-1732.
34
35
36 27. Cauchi, M. R.; Henchal, E.; Wright, P. J., The sensitivity of cell-associated dengue virus proteins
37 to trypsin and the detection of trypsin-resistant fragments of the nonstructural glycoprotein NS1.
38 *Virology* **1991**, 180, (2), 659-667.
39
40
41 28. Bezouška, K.; Sklenář, J.; Novák, P.; Halada, P.; Havlíček, V.; Kraus, M.; Tichá, M.; Jonáková, V.,
42 Determination of the complete covalent structure of the major glycoform of DQH sperm surface protein,
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 a novel trypsin-resistant boar seminal plasma O-glycoprotein related to pB1 protein. *Protein science*

4
5
6 **1999**, 8, (7), 1551-1556.

7
8 29. Takahashi, K.; Smith, A. D.; Poulsen, K.; Kilian, M.; Julian, B. A.; Mestecky, J.; Novak, J.; Renfrow,

9
10 M. B., Naturally Occurring Structural Isomers in Serum IgA1 O-Glycosylation. *Journal of Proteome*

11
12 *Research* **2011**, 11, (2), 692-702.

13
14 30. Wada, Y.; Dell, A.; Haslam, S. M.; Tissot, B.; Canis, K.; Azadi, P.; Bäckström, M.; Costello, C. E.;

15
16 Hansson, G. C.; Hiki, Y., Comparison of Methods for Profiling O-Glycosylation HUMAN PROTEOME

17
18 ORGANISATION HUMAN DISEASE GLYCOMICS/PROTEOME INITIATIVE MULTI-INSTITUTIONAL STUDY OF

19
20 IgA1. *Molecular & Cellular Proteomics* **2010**, 9, (4), 719-727.

21
22 31. Renfrow, M. B.; Mackay, C. L.; Chalmers, M. J.; Julian, B. A.; Mestecky, J.; Kilian, M.; Poulsen, K.;

23
24 Emmett, M. R.; Marshall, A. G.; Novak, J., Analysis of O-glycan heterogeneity in IgA1 myeloma proteins

25
26 by Fourier transform ion cyclotron resonance mass spectrometry: implications for IgA nephropathy.

27
28 *Analytical and bioanalytical chemistry* **2007**, 389, (5), 1397-1407.

29
30 32. Renfrow, M. B.; Cooper, H. J.; Tomana, M.; Kulhavy, R.; Hiki, Y.; Toma, K.; Emmett, M. R.;

31
32 Mestecky, J.; Marshall, A. G.; Novak, J., Determination of aberrant O-glycosylation in the IgA1 hinge

33
34 region by electron capture dissociation fourier transform-ion cyclotron resonance mass spectrometry.

35
36 *Journal of Biological Chemistry* **2005**, 280, (19), 19136-19145.

37
38 33. Takahashi, K.; Wall, S. B.; Suzuki, H.; Smith, A. D.; Hall, S.; Poulsen, K.; Kilian, M.; Mobley, J. A.;

39
40 Julian, B. A.; Mestecky, J.; Novak, J.; Renfrow, M. B., Clustered O-Glycans of IgA1: DEFINING MACRO-

41
42 AND MICROHETEROGENEITY BY USE OF ELECTRON CAPTURE/TRANSFER DISSOCIATION. *Molecular &*

43
44 *Cellular Proteomics* **2010**, 9, (11), 2545-2557.

45
46 34. Seipert, R. R.; Dodds, E. D.; Lebrilla, C. B., Exploiting differential dissociation chemistries of O-

47
48 linked glycopeptide ions for the localization of mucin-type protein glycosylation. *Journal of proteome*

49
50 *research* **2008**, 8, (2), 493-501.

51
52
53
54
55
56
57
58
59
60

- 1
2
3 35. Seipert, R. R.; Dodds, E. D.; Clowers, B. H.; Beecroft, S. M.; German, J. B.; Lebrilla, C. B., Factors
4 that influence fragmentation behavior of N-linked glycopeptide ions. *Analytical chemistry* **2008**, *80*, (10),
5
6 3684-3692.
7
8
9
10 36. Nwosu, C. C.; Seipert, R. R.; Strum, J. S.; Hua, S. S.; An, H. J.; Zivkovic, A. M.; German, B. J.;
11
12 Lebrilla, C. B., Simultaneous and extensive site-specific N-and O-glycosylation analysis in protein
13
14 mixtures. *Journal of proteome research* **2011**, *10*, (5), 2612-2624.
15
16
17 37. Hua, S.; Nwosu, C. C.; Strum, J. S.; Seipert, R. R.; An, H. J.; Zivkovic, A. M.; German, J. B.; Lebrilla,
18
19 C. B., Site-specific protein glycosylation analysis with glycan isomer differentiation. *Analytical and*
20
21 *bioanalytical chemistry* **2012**, *403*, (5), 1291-1302.
22
23
24 38. Clowers, B. H.; Dodds, E. D.; Seipert, R. R.; Lebrilla, C. B., Site determination of protein
25
26 glycosylation based on digestion with immobilized nonspecific proteases and Fourier transform ion
27
28 cyclotron resonance mass spectrometry. *Journal of proteome research* **2007**, *6*, (10), 4032-4040.
29
30
31 39. Strum, J. S.; Nwosu, C. C.; Hua, S.; Kronewitter, S. R.; Seipert, R. R.; Bachelor, R. J.; An, H. J.;
32
33 Lebrilla, C. B., Automated Assignments of N-and O-Site Specific Glycosylation with Extensive Glycan
34
35 Heterogeneity of Glycoprotein Mixtures. *Analytical chemistry* **2013**, *85*, (12), 5666-5675.
36
37
38 40. Hong, Q.; Lebrilla, C. B.; Miyamoto, S.; Ruhaak, L. R., Absolute quantitation of immunoglobulin G
39
40 and its glycoforms using multiple reaction monitoring. *Analytical chemistry* **2013**, *85*, (18), 8585-8593.
41
42
43 41. Langley, R.; Wines, B.; Willoughby, N.; Basu, I.; Proft, T.; Fraser, J. D., The staphylococcal
44
45 superantigen-like protein 7 binds IgA and complement C5 and inhibits IgA-Fc α RI binding and serum
46
47 killing of bacteria. *The Journal of Immunology* **2005**, *174*, (5), 2926-2933.
48
49
50 42. Wada, Y.; Tajiri, M.; Yoshida, S., Hydrophilic affinity isolation and MALDI multiple-stage tandem
51
52 mass spectrometry of glycopeptides for glycoproteomics. *Analytical chemistry* **2004**, *76*, (22), 6560-6565.
53
54
55 43. Ninonuevo, M.; An, H.; Yin, H.; Killeen, K.; Grimm, R.; Ward, R.; German, B.; Lebrilla, C.,
56
57 Nanoliquid chromatography-mass spectrometry of oligosaccharides employing graphitized carbon
58
59
60

1
2
3 chromatography on microchip with a high-accuracy mass analyzer. *Electrophoresis* **2005**, *26*, (19), 3641-
4
5 3649.

6
7
8 44. Chu, C. S.; Niñonuevo, M. R.; Clowers, B. H.; Perkins, P. D.; An, H. J.; Yin, H.; Killeen, K.; Miyamoto,
9
10 S.; Grimm, R.; Lebrilla, C. B., Profile of native N-linked glycan structures from human serum using high
11
12 performance liquid chromatography on a microfluidic chip and time-of-flight mass spectrometry.
13
14 *Proteomics* **2009**, *9*, (7), 1939-1951.

15
16
17 45. Nwosu, C. C.; Seipert, R. R.; Strum, J. S.; Hua, S. S.; An, H. J.; Zivkovic, A. M.; German, B. J.;
18
19 Lebrilla, C. B., Simultaneous and Extensive Site-specific N- and O-Glycosylation Analysis in Protein
20
21 Mixtures. *Journal of Proteome Research* **2011**, *10*, (5), 2612-2624.

22
23
24 46. Picariello, G.; Ferranti, P.; Mamone, G.; Roepstorff, P.; Addeo, F., Identification of N-linked
25
26 glycoproteins in human milk by hydrophilic interaction liquid chromatography and mass spectrometry.
27
28 *Proteomics* **2008**, *8*, (18), 3833-3847.

29
30
31 47. Hua, S.; Hu, C. Y.; Kim, B. J.; Totten, S. M.; Oh, M. J.; Yun, N.; Nwosu, C. C.; Yoo, J. S.; Lebrilla, C.
32
33 B.; An, H. J., Glyco-Analytical Multispecific Proteolysis (Glyco-AMP): A Simple Method for Detailed and
34
35 Quantitative Glycoproteomic Characterization. *Journal of Proteome Research* **2013**, *12*, (10), 4414-4423.

36
37
38 48. Krugmann, S.; Pleass, R.; Atkin, J.; Woof, J., Mutagenesis of J chain residues critical for IgA dimer
39
40 assembly. *Biochemical Society transactions* **1997**, *25*, (2), 323S.

41
42
43 49. Vaerman, J.-P.; Langendries, A.; Giffroy, D.; Brandtzaeg, P.; Kobayashi, K., Lack of SC/pIgR-
44
45 mediated epithelial transport of a human polymeric IgA devoid of J chain: in vitro and in vivo studies.
46
47 *Immunology* **1998**, *95*, (1), 90.

48
49
50 50. Mattu, T. S.; Pleass, R. J.; Willis, A. C.; Kilian, M.; Wormald, M. R.; Lellouch, A. C.; Rudd, P. M.;
51
52 Woof, J. M.; Dwek, R. A., The glycosylation and structure of human serum IgA1, Fab, and Fc regions and
53
54 the role of N-glycosylation on Fc α receptor interactions. *Journal of Biological Chemistry* **1998**, *273*, (4),
55
56 2260-2272.

Figure Legends:

Figure 1. Extracted Compound chromatogram (ECC) and associated structural assignments of glycopeptides from pronase digested sIgA isolated from an individual donor. (A) ECC of major glycopeptides eluted between 6 and 38 min; (B) ECC of all glycopeptides eluted between 7 and 21 min; (C) ECC of all glycopeptides eluted between 21 to 29 min. Glycosites corresponding to the four polypeptides was denoted in different colors. Green Circles, yellow circles, blue squares, red triangles, and purple diamonds represent mannose, galactose, GlcNAc, fucose and Neu5Ac, respectively.

Figure 2. Deconvoluted MS/MS spectra of three N-linked glycopeptides from sIgA. (A) MS/MS data for a N-linked peptide from J chain; (B) MS/MS data for a neutral N-linked peptide from SC; (C) MS/MS data for an acidic N-linked peptide from SC. Green Circles, yellow circles, blue squares, red triangles, and purple diamonds represent mannose, galactose, GlcNAc, fucose and Neu5Ac, respectively.

Figure 3. Extracted compound chromatogram (ECC) of glycopeptides from pronase digested commercial pooled human colostrum IgA with corresponding structural assignments. Different colors represent various sites. Green Circles, yellow circles, blue squares, red triangles, and purple diamonds represent mannose, galactose, GlcNAc, fucose and Neu5Ac, respectively.

Figure 4. (A-C) Glycosite occupancy and heterogeneity in pronase digested sIgA isolated from an individual donor, SC, J chain, and IgA1/IgA2, respectively. (D-F) Glycosite occupancy and heterogeneity in pooled commercial sIgA, SC, J chain, and IgA1/IgA2, respectively. Circle shading under each site is representative of site occupancy ratio to the most occupied site in

1
2
3 ^{92}Asn from IgA2. Circle shading under each glycan is representative of the relative glycan
4 abundance at each site.
5
6
7
8
9

10 **Figure 5.** MS spectra of three tryptic O-glycopeptides from sIgA and the corresponding isotopic
11 patterns and tandem spectra. (A) O-glycopeptide with glycan composition
12 HexNAc₅Hex₄Neu5Ac₃; (B) O-glycopeptide with glycan composition HexNAc₅Hex₄Neu5Ac₂;
13 (C) O-glycopeptide with glycan composition HexNAc₅Hex₄. Red boxes are representative of the
14 predicted isotopic pattern.
15
16
17
18
19
20
21
22
23
24

25 **Figure 6:** Extracted compound chromatograms (ECCs) of four sIgA HR O-glycopeptides
26 containing same number of N-acetylgalactosamine and galactose residues and 0-3 sialic acid
27 monosaccharides.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1.




Figure 2.




Figure 3.





Figure 4.






Figure 6.




Table 1. List of N-glycopeptides from isolated sIgA from collected milk sample

RT (min)	M/Z	Mass (Da)	PPM error	Protein	Sequence	Site	Hex	HexNAc	Fuc	Neu5Ac	Intensity
33.869	1187.807	3560.4003	5	IGA2	HVKHYTNPSQ	92	5	4	1	2	68197072
13.293	832.324	2493.9477	8	IGA2	YTNPS	92	5	4	0	1	464028
26.164	878.6541	2632.9404	20	IGA2	YTNPSQDV	92	5	3	0	1	775421
26.196	946.348	2836.0191	20	IGA2	YTNPSQDV	92	5	4	0	1	2130024
29.815	1043.384	3127.1324	13	IGA2	YTNPSQDV	92	5	4	0	2	3291937
23.32	977.0724	2928.1911	9	IGJ	ENISDPTSPL	71	5	3	1	1	1381861
26.32	1141.784	3422.3268	3	IGJ	ENISDPTSPL	71	5	4	1	2	1661523
21.692	953.0611	2856.1589	5	IGJ	NISDPTSPL	71	5	4	0	1	860383
22.502	1001.749	3002.2246	7	IGJ	NISDPTSPL	71	5	4	1	1	329314
25.502	1050.09	3147.2477	2	IGJ	NISDPTSPL	71	5	4	0	2	1405679
17.758	900.6658	2698.9762	6	SC	NDT	135	5	4	1	2	777521
24.384	1147.841	3440.5043	0	SC	VSLEVSQGPGLLNDTK	135	6	4	0	0	1123577
24.136	1215.534	3643.5889	0	SC	VSLEVSQGPGLLNDTK	135	6	5	0	0	4305616
20.277	952.4099	3805.6105	7	SC	VSLEVSQGPGLLNDTK	135	7	5	0	0	253514
23.925	952.4168	3805.6334	1	SC	VSLEVSQGPGLLNDTK	135	7	5	0	0	6387259
23.798	962.6741	3846.6648	0	SC	VSLEVSQGPGLLNDTK	135	6	6	0	0	4180562
19.877	1003.179	4008.685	8	SC	VSLEVSQGPGLLNDTK	135	7	6	0	0	174310
23.585	1003.189	4008.7254	1	SC	VSLEVSQGPGLLNDTK	135	7	6	0	0	7677994
20.071	847.3634	3385.4165	7	SC	NYTGRIRLD	186	6	5	0	1	1881945
19.575	898.1307	3588.496	6	SC	NYTGRIRLD	186	6	6	0	1	1677348

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

16.882	850.3206	2547.9384	6	SC	NPN/PNGG	186/421	5	4	0	2	1504821
18.096	815.9875	2444.9406	3	SC	VPGN(PGNV)	469	5	4	1	1	508112
20.296	913.0201	2736.0374	3	SC	VPGN(PGNV)	469	5	4	1	2	1157752
24.476	1020.688	4078.721	13	SC	LTNFPENGTFFVNNIAQLSQ	90	6	5	0	0	6116531
26.152	1052.947	4207.7536	15	SC	LTNFPENGTFFVNNIAQLSQ	90	5	5	0	1	3595040
24.225	1071.457	4281.7958	14	SC	LTNFPENGTFFVNNIAQLSQ	90	6	6	0	0	6857053
26.003	1093.458	4369.8037	15	SC	LTNFPENGTFFVNNIAQLSQ	90	6	5	0	1	8980128
25.819	1103.718	4410.835	14	SC	LTNFPENGTFFVNNIAQLSQ	90	5	6	0	1	4215505
24.654	1139.731	4554.8928	10	SC	LTNFPENGTFFVNNIAQLSQ	90	6	6	0	1	3272015
25.691	1144.232	4572.8955	12	SC	LTNFPENGTFFVNNIAQLSQ	90	6	6	0	1	9554195
26.363	1161.731	4642.8965	13	SC	LTNFPENGTFFVNNIAQLSQ	90	6	5	0	2	4439261
25.359	1184.75	4734.9699	7	SC	LTNFPENGTFFVNNIAQLSQ	90	7	6	0	1	3199461
26.062	1212.498	4845.959	16	SC	LTNFPENGTFFVNNIAQLSQ	90	6	6	0	2	1359938
24.086	1212.501	4845.9763	12	SC	LTNFPENGTFFVNNIAQLSQ	90	6	6	0	2	597825
27.84	1217.005	4863.9906	11	SC	LTNFPENGTFFVNNIAQLSQ	90	6	6	0	2	6714054
25.776	984.6771	3934.6744	9	SC	EGYVSSKYAGRANLT	83	5	6	0	1	4837727
25.585	1025.191	4096.7328	10	SC	EGYVSSKYAGRANLT	83	6	6	0	1	9760395
22.825	1025.194	4096.7449	13	SC	EGYVSSKYAGRANLT	83	6	6	0	1	990066
25.424	1035.447	4137.7591	10	SC	EGYVSSKYAGRANLT	83	5	7	0	1	5184296
22.162	1075.941	4299.7436	5	SC	EGYVSSKYAGRANLT	83	6	7	0	1	924140
25.21	1075.962	4299.8158	11	SC	EGYVSSKYAGRANLT	83	6	7	0	1	12223087
28.002	1097.964	4387.8253	9	SC	EGYVSSKYAGRANLT	83	6	6	0	2	6682655
24.758	1097.964	4387.8259	9	SC	EGYVSSKYAGRANLT	83	6	6	0	2	1611696
27.447	1148.735	4590.9081	9	SC	EGYVSSKYAGRANLT	83	6	7	0	2	8800010
9.058	710.7651	1419.5156	1	SC//IGA1//IGA2	AN	83//144//131/205	5	2	0	0	266033
8.906	617.9116	1850.7109	3	SC//IGA1//IGA2	AN	83//144//131/205	3	5	1	0	548567
9.366	671.9295	2012.7634	3	SC//IGA1//IGA2	AN	83//144//131/205	4	5	1	0	361962
9.703	720.275	2157.8008	3	SC//IGA1//IGA2	AN	83//144//131/205	4	5	0	1	598056
10.14	774.292	2319.8541	3	SC//IGA1//IGA2	AN	83//144//131/205	5	5	0	1	343641
15.706	822.9688	2465.8852	8	SC//IGA1//IGA2	AN	83//144//131/205	5	5	1	1	199925
12.099	822.9805	2465.9174	5	SC//IGA1//IGA2	AN	83//144//131/205	5	5	1	1	261020
9.281	961.37	1920.725	2	SC//IGA1//IGA2	ENGT//SEAN	90//144//131	3	5	0	0	493651
10.416	656.2664	1965.7749	20/14//4	SC//IGA1//IGA2	ENG/WN//NVS	90/499//340/327	3	5	1	0	556088
10.699	710.2852	2127.8318	20/15//5	SC//IGA1//IGA2	ENG/WN//NVS	90/499//340/327	4	5	1	0	1079828
13.368	807.3168	2418.9288	20/15//5	SC//IGA1//IGA2	ENG/WN//NVS	90/499//340/327	4	5	1	1	640041
13.258	861.3339	2580.9775	17/11//3	SC//IGA1//IGA2	ENG/WN//NVS	90/499//340/327	5	5	1	1	715784
16.21	958.3694	2872.0847	19/14//7	SC//IGA1//IGA2	ENG/WN//NVS	90/499//340/327	5	5	1	2	610683
11.434	724.7811	1447.5476	1	IGA1//IGA2//SC	NV	340//47/327//469	5	2	0	0	254065
9.941	805.8059	1609.5973	0	IGA1//IGA2//SC	NV	340//47/327//469	6	2	0	0	153825
10.484	681.271	2040.7925	2	IGA1//IGA2//SC	NV	340//47/327//469	4	5	1	0	582738

8.54	917.8562	1833.6979	5//14//14	IGA1//IGA2//SC//IGJ	EAN//NVT//NIS	144//47//131//469//71	3	5	0	0	325261
19.811	921.6835	2762.0284	22//4	IGA1//IGA2//SC	LLGSEAN//EPPGNGT	131//144//421	5	4	1	1	586605
19.407	970.3717	2908.0924	19//6	IGA1//IGA2//SC	LLGSEAN//EPPGNGT	131//144//421	5	4	2	1	306195
22.092	1018.719	3053.1319	17//6	IGA1//IGA2//SC	LLGSEAN//EPPGNGT	131//144//421	5	4	1	2	1448849

Table 2. List of sIgA HR O-glycopeptides

RT (min)	Base Peak (Da)	Z	Error (ppm)	Formula	Hex	HexNAc	Neu5Ac	Relative intensity (%)
22.35	1441.7946	5	-3.5	C294 H459 N59 O143 S3	6	6	3	0.07
22.12	1383.5586	5	-15.9	C283 H442 N58 O135 S3	6	6	2	0.07
21.93	1325.1411	5	-15.4	C272 H425 N57 O127 S3	6	6	1	0.04
23.12	1467.5875	5	-14.1	C299 H466 N60 O146 S3	6	5	4	0.15
22.55	1409.3685	5	-14.6	C288 H449 N59 O138 S3	6	5	3	0.45
22.28	1351.1494	5	-15.3	C277 H432 N58 O130 S3	6	5	2	0.57
22.04	1292.9313	5	-15.2	C266 H415 N57 O122 S3	6	5	1	0.32
21.76	1234.7120	5	-16.1	C255 H398 N56 O114 S3	6	5	0	0.08
23.40	1435.1755	5	-15.4	C293 H456 N60 O141 S3	6	4	4	0.38
22.83	1376.9572	5	-15.5	C282 H439 N59 O133 S3	6	4	3	1.64
22.49	1318.7403	5	-14.6	C271 H422 N58 O125 S3	6	4	2	1.35
22.25	1260.1283	5	-9.6	C260 H405 N57 O117 S3	6	4	1	0.31
21.95	1502.3845	4	-10.2	C249 H388 N56 O109 S3	6	4	0	0.14
23.59	1344.5323	5	-26.5	C276 H429 N59 O128 S3	6	3	3	0.94
23.20	1286.3315	5	-13.5	C265 H412 N58 O120 S3	6	3	2	1.15
22.81	1227.9189	5	-8.9	C254 H395 N57 O112 S3	6	3	1	0.70
22.43	1169.7008	5	-8.5	C243 H378 N56 O104 S3	6	3	0	0.34
24.71	1485.1758	5	-24.0	C302 H470 N60 O149 S3	5	5	5	0.59
23.72	1427.1532	5	-27.4	C291 H453 N59 O141 S3	5	5	4	1.87
23.19	1368.7517	5	-15.7	C280 H436 N58 O133 S3	5	5	3	2.39
22.78	1310.5358	5	-14.0	C269 H419 N57 O125 S3	5	5	2	2.21
22.39	1252.3178	5	-13.8	C258 H402 N56 O117 S3	5	5	1	1.56
22.10	1193.9069	5	-7.6	C247 H385 N55 O109 S3	5	5	0	0.57
24.56	1452.7646	5	-24.9	C296 H460 N60 O144 S3	5	4	5	0.10
24.18	1394.5549	5	-19.2	C285 H443 N59 O136 S3	5	4	4	6.72
23.69	1336.3333	5	-21.9	C274 H426 N58 O128 S3	5	4	3	9.20
23.13	1278.1190	5	-19.2	C263 H409 N57 O120 S3	5	4	2	7.93
22.65	1219.7091	5	-12.6	C252 H392 N56 O112 S3	5	4	1	5.30
22.29	1161.4932	5	-10.5	C241 H375 N55 O104 S3	5	4	0	1.85
24.41	1303.9263	5	-19.8	C268 H416 N58 O123 S3	5	3	3	3.78
23.93	1245.5141	5	-15.2	C257 H399 N57 O115 S3	5	3	2	4.08
23.27	1187.3040	5	-8.4	C246 H382 N56 O107 S3	5	3	1	2.48
22.76	1129.0857	5	-8.1	C235 H365 N55 O99 S3	5	3	0	0.68
24.79	1213.1057	5	-13.8	C251 H389 N57 O110 S3	5	2	2	0.33

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

24.10	1154.8932	5	-8.8	C240 H372 N56 O102 S3	5	2	1	0.37
25.42	1412.1573	5	-19.6	C288 H447 N59 O139 S3	4	4	5	0.27
25.14	1353.9268	5	-28.9	C277 H430 N58 O131 S3	4	4	4	6.64
24.74	1295.7254	5	-16.5	C266 H413 N57 O123 S3	4	4	3	6.67
24.14	1237.5065	5	-17.1	C255 H396 N56 O115 S3	4	4	2	5.48
23.46	1179.0949	5	-11.6	C244 H379 N55 O107 S3	4	4	1	3.20
22.87	1120.8728	5	-15.0	C233 H362 N54 O99 S3	4	4	0	0.96
25.65	1321.5394	5	-12.0	C271 H420 N58 O126 S3	4	3	4	0.31
25.54	1263.3206	5	-12.4	C260 H403 N57 O118 S3	4	3	3	4.12
25.13	1204.8921	5	-20.7	C249 H386 N56 O110 S3	4	3	2	3.67
24.34	1146.6773	5	-18.1	C238 H369 N55 O102 S3	4	3	1	1.63
23.56	1088.4603	5	-17.2	C227 H352 N54 O94 S3	4	3	0	0.50
26.05	977.2487	6	-8.2	C243 H376 N56 O105 S3	4	2	2	0.16
24.69	928.7259	6	-16.0	C232 H359 N55 O97 S3	4	2	1	0.06
25.75	1222.5168	5	-2.9	C252 H390 N56 O113 S3	3	3	3	2.60
25.16	1164.2785	5	-19.5	C241 H373 N55 O105 S3	3	3	2	1.62
24.18	1106.2704	5	-10.7	C230 H356 N54 O97 S3	3	3	1	0.24
26.16	1131.8795	5	-9.9	C235 H363 N55 O100 S3	3	2	2	1.09
28.19	1199.5245	4	-8.6	C202 H310 N52 O77 S3	1	1	1	0.06
28.80	1035.4722	4	-5.6	C177 H270 N50 O59 S3	0	0	0	0.02

